arXiv:1701.08690v1 [math.AG] 30 Jan 2017

ON THE EXISTENCE OF UNIVERSAL FAMILIES OF MARKED HYPERKÄHLER VARIETIES

EYAL MARKMAN

Abstract. We prove the existence of a universal family over every component of the moduli space of marked irreducible holomorphic symplectic manifolds. The analogous result follows for the Teichmüller spaces.

Contents

1. Introduction 1
2. Moduli spaces of marked pairs and triples 3
3. Gerbes 6
4. Vanishing of the class of a gerbe 8
References 10

1. Introduction

An *irreducible holomorphic symplectic manifold* is a simply connected compact Kähler manifold X, such that $H^0(X,\Omega_X^2)$ is one dimensional spanned by an everywhere non-degenerate holomorphic 2-form [Be1, Hu1]. The second integral cohomology of X is endowed with a non-degenerate integral symmetric bilinear pairing of signature $(3,b_2-3)$, where b_2 is the second Betti number of X. The pairing is known as the *Beauville-Bogomolov-Fujiki* pairing. A *marking* for X is an isometry $\eta: H^2(X,\mathbb{Z})\to\Lambda$ with a fixed lattice Λ. An isomorphism of two marked pairs (X_i,η_i), $i=1,2$, consists of an isomorphism $f: X_1\to X_2$, such that $\eta_1\circ f^*=\eta_2$.

Given an analytic space B and a discrete group H, denote by $\underline{H}_B$ the trivial local system over B with fiber H. A *family of marked irreducible holomorphic symplectic manifolds* over an analytic base B consists of a family $\pi:\mathcal{X}\to B$ of such manifolds, together with an isometric trivialization $\eta: R^2\pi_*\mathbb{Z}\to\underline{\Lambda}_B$. We will call the pair (π,η) a Λ-*marked* family for short. Two Λ-marked families $(\pi:\mathcal{X}\to B,\eta)$ and $(\tilde\pi:\tilde{\mathcal{X}}\to B,\tilde\eta)$ are *isomorphic*, if there exists an isomorphism $f:\mathcal{X}\to\tilde{\mathcal{X}}$, such that $\tilde\pi f=\pi$ and $\tilde\eta=\eta\circ f^*$, where $f^*: R^2\tilde\pi_*\mathbb{Z}\to R^2\pi_*\mathbb{Z}$ is the isomorphism induced by f. Given a marked family $(\pi:\mathcal{X}\to B,\eta)$ and a morphism $\kappa:\tilde B\to B$, we get the pulled back family $\kappa^*(\pi):\mathcal{X}\times_B\tilde B\to\tilde B$ with the marking $\kappa^*(\eta)$. Let $\mathcal{F}_\Lambda$ be the functor, from the category of analytic spaces to the category of sets, which associates to an analytic space B the set of isomorphism classes of Λ-marked families (π,η) over B. There exists a non-Hausdorff (disconnected) complex manifold $\mathfrak{M}_\Lambda$ of dimension $\mathrm{rank}(\Lambda)-2$, which coarsely represents $\mathcal{F}_\Lambda$ [Hu1].

Let X be an irreducible holomorphic symplectic manifold. Denote by $\mathrm{Aut}_0(X)$ the subgroup, of the auotomorphism group $\mathrm{Aut}(X)$ of X, consisting of elements which act trivially on

Date: January 31, 2017.

$H^2(X,\mathbb{Z})$. Huybrechts proved that $\mathrm{Aut}_0(X)$ is a finite group [Hu1, Prop. 9.1(v)]. Associated to every family $\pi : \mathcal{X} \to B$ of irreducible holomorphic symplectic manifolds we have a local system $\mathcal{A}ut_0(\pi)$ over B, whose fiber over $b \in B$ is the group $\mathrm{Aut}_0(X_b)$ of the fiber X_b of π over b, by [HT, Theorem 2.1].

Given an $\mathcal{A}ut_0(\pi)$ torsor $\mathcal{P}$ over B, we get the family $\tilde{\mathcal{X}} := \mathcal{X} \times^{\mathcal{A}ut_0(\pi)} \mathcal{P}$. Denote by $\tilde{\pi} : \tilde{\mathcal{X}} \to B$ the natural projection. Note that the local systems $R^2\pi_*\mathbb{Z}$ and $R^2\tilde{\pi}_*\mathbb{Z}$ are naturally isomorphic, and so a marking η for the former induces a marking $\tilde{\eta}$ for the latter. We denote this new marked family by $(\tilde{\pi}, \tilde{\eta}) := (\pi, \eta) \times \mathcal{P}$. Two Λ-marked families $(\pi : \mathcal{X} \to B, \eta)$ and $(\tilde{\pi} : \tilde{\mathcal{X}} \to B, \tilde{\eta})$ are said to be *equivalent*, if there exists an $\mathcal{A}ut_0(\pi)$ torsor $\mathcal{P}$ over B, such that $(\tilde{\pi}, \tilde{\eta}) := (\pi, \eta) \times \mathcal{P}$. The map $\mathcal{P} \mapsto (\pi, \eta) \times \mathcal{P}$ is a bijection between the set $H^1(B, \mathcal{A}ut_0(\pi))$, of isomorphism classes of $\mathcal{A}ut_0(\pi)$ torsors, and the set of isomorphism classes of Λ-marked families equivalent to (π, η). Its inverse sends $(\tilde{\pi}, \tilde{\eta})$ to the isomorphism class of the $\mathcal{A}ut_0(\pi)$ torsor $\mathcal{I}som((\pi, \eta), (\tilde{\pi}, \tilde{\eta}))$ of local isomorphisms of the two families compatible with the markings. We elaborate on this bijection in Remark 4.4.

Set $\underline{\Lambda} := \underline{\Lambda}_{\mathfrak{M}_\Lambda}$. The main result of this paper is the following statement.

Theorem 1.1. *There exists a family $\pi : \mathcal{X} \to \mathfrak{M}_\Lambda$ of irreducible holomorphic symplectic manifolds and a marking $\eta : R^2\pi_*\mathbb{Z} \to \underline{\Lambda}$ satisfying the following universal property. Given a family of Λ-marked irreducible holomorphic symplectic manifolds $(\tilde{\pi} : \tilde{\mathcal{X}} \to B, \tilde{\eta})$ over an analytic space B, the pullback $(\kappa^*(\pi), \kappa^*(\eta))$ via the classifying morphism $\kappa : B \to \mathfrak{M}_\Lambda$ is equivalent to $(\tilde{\pi}, \tilde{\eta})$. The marked family (π, η) satisfying this property is unique, up to isomorphism. $\mathcal{A}ut_0(\pi)$ restricts to each connected component of $\mathfrak{M}_\Lambda$ as a trivial local system.*

The Theorem is proved in section 4. The Teichmüller space of an irreducible holomorphic symplectic manifold X maps to the moduli space of marked pairs [Ver, Cor. 4.31], and so the universal family over the latter pulls back to one over the Teichmüller space.

Consider the moduli space $\mathfrak{M}_{\Lambda,G}$ of isomorphism classes of triples (X, η, ψ), where (X, η) is a marked irreducible holomorphic symplectic manifold, and $\psi : \mathrm{Aut}_0(X) \to G$ is an isomorphism with a fixed finite group G. The automorphism group $\mathrm{Aut}(X, \eta)$ of the marked pair (X, η) is $\mathrm{Aut}_0(X)$, and the automorphism group $\mathrm{Aut}(X, \eta, \psi)$ of the triple is the center of $\mathrm{Aut}_0(X)$.

Remark 1.2. If the center of G is trivial, then the moduli space $\mathfrak{M}_{\Lambda,G}$ represents the functor from the category of analytic spaces to the category of sets, which associates to an analytic space B the set of isomorphism classes of triples (π, η, ψ), consisting of a family $\pi : \mathcal{X} \to B$ of irreducible holomorphic symplectic manifolds, an isometric trivialization $\eta : R^2\pi_*\mathbb{Z} \to \underline{\Lambda}_B$, and a trivialization $\psi : \mathcal{A}ut_0(\pi) \to \underline{G}_B$.

Let us sketch the proof of Theorem 1.1. We have the forgetful morphism $\phi : \mathfrak{M}_{\Lambda,G} \to \mathfrak{M}_\Lambda$. We first show that ϕ restricts to each connected component $\mathfrak{M}^0_{\Lambda,G}$ of $\mathfrak{M}_{\Lambda,G}$ as an isomorphism onto the corresponding connected component $\mathfrak{M}^0_\Lambda$ of $\mathfrak{M}_\Lambda$ (Lemma 2.2).

Fix a connected component $\mathfrak{M}^0_{\Lambda,G}$ of $\mathfrak{M}_{\Lambda,G}$. Let $\underline{\Lambda}$ be the trivial local system over $\mathfrak{M}^0_{\Lambda,G}$ with fiber Λ and define $\underline{G}$ similarly. Consider the stack $\mathbb{G}$ over $\mathfrak{M}^0_{\Lambda,G}$, which associates to each subset U of $\mathfrak{M}^0_{\Lambda,G}$, open in the classical topology, the following category $\mathbb{G}(U)$. Objects of $\mathbb{G}(U)$ are triples (π, η, ψ), consisting of a family $\pi : \mathcal{X} \to U$ of irreducible holomorphic symplectic manifolds, an isometric isomorphism of local systems $\eta : R^2\pi_*\mathbb{Z} \to \underline{\Lambda}_U$, and an isomorphism of local systems $\psi : \mathcal{A}ut_0(\pi) \to \underline{G}_U$, such that the triple (X_t, η_t, ψ_t) over a point t of U represents the isomorphism class parametrized by t as a point of the coarse moduli space $\mathfrak{M}^0_{\Lambda,G}$. The morphisms of $\mathbb{G}(U)$ are isomorphisms of families, compatible with the trivializations of the two local systems.

We observe next that $\mathbb{G}$ is a gerbe over $\mathfrak{M}^0_{\Lambda,G}$ with band the trivial sheaf of groups $\underline{Z}$ with fiber the center Z of G (Lemma 3.4). See section 3 for the definitions. Equivalence classes of gerbes with band $\underline{Z}$ are parametrized by $\check{H}^2(\mathfrak{M}^0_{\Lambda,G}, Z)$ [Gi, Br, Theorem 5.2.9]. The existence of a universal family over $\mathfrak{M}^0_{\Lambda,G}$ is equivalent to the vanishing of the class $[\mathbb{G}] \in \check{H}^2(\mathfrak{M}^0_{\Lambda,G}, Z)$. Denote by $\phi_*[\mathbb{G}]$ the image of $[\mathbb{G}]$ in $\check{H}^2(\mathfrak{M}^0_{\Lambda}, Z)$ via the isomorphism $\phi : \mathfrak{M}^0_{\Lambda,G} \to \mathfrak{M}^0_{\Lambda}$.

Associated to a Kähler class ω on X and a marking η is a *twistor line* $Tw_{\omega,\eta}$ in $\mathfrak{M}^0_{\Lambda}$ through the marked pair (X, η) [Hu1, 1.13]. $Tw_{\omega,\eta}$ is isomorphic to a projective line $\mathbb{P}^1$. Finally, we show that the restriction homomorphism $\check{H}^2(\mathfrak{M}^0_{\Lambda}, Z) \to \check{H}^2(Tw_{\omega,\eta}, Z)$ is an isomorphism (Lemma 4.2), using the Global Torelli Theorem [Ver]. The image of $\phi_*[\mathbb{G}]$ in $\check{H}^2(Tw_{\omega,\eta}, Z)$ vanishes, by the existence of the twistor family over $Tw_{\omega,\eta}$. Hence, the class $[\mathbb{G}]$ is trivial and Theorem 1.1 follows.

Irreducible holomorphic symplectic manifolds of $K3^{[n]}$-*type* are those, which are deformation equivalent to the Hilbert scheme of length n subschemes of a $K3$ surface. If X is of $K3^{[n]}$-type, then any automorphism of X, which acts trivially on $H^2(X, \mathbb{Z})$, is the identity. This follows for Hilbert schemes by a result of Beauville [Be2], and consequently also for their deformations, see [HT, Sec. 2]. The automorphism group of every marked pair (X, η), with X of $K3^{[n]}$-type, is thus trivial, and Theorem 1.1 is known in this case.

Example 1.3. Fix an integer $n \geq 2$. Let T be a two-dimensional compact complex torus, $T^{[n+1]}$ its Douady space of length $n+1$ subschemes, $T^{(n+1)}$ its $(n+1)$ symmetric product, and consider the fiber $K^{[n]}(T)$ over $0 \in T$ of the composition $T^{[n+1]} \to T^{(n+1)} \to T$, where the left arrow is the Hilbert Chow morphism, and the right is summation. $K^{[n]}(T)$ is a $2n$-dimensional irreducible holomorphic symplectic manifold known as the *generalized Kummer manifold* associated to T [Be1]. Translation by points of T of order $n+1$ induce automorphisms of $T^{[n+1]}$ leaving $K^{[n]}(T)$ invariant, as does multiplication by -1. These automorphisms generate $\mathrm{Aut}_0(K^{[n]}(T))$. When X is deformation equivalent to $K^{[n]}(T)$, $\mathrm{Aut}_0(X)$ is thus isomorphic to the semidirect product $[\mathbb{Z}/(n+1)\mathbb{Z}]^4 \rtimes \mathbb{Z}/2\mathbb{Z}$, where the non-trivial element of $\mathbb{Z}/2\mathbb{Z}$ acts on $[\mathbb{Z}/(n+1)\mathbb{Z}]^4$ via multiplication by -1 (see [BSN, Theorem 3 and Corollary 5]). The center of $\mathrm{Aut}_0(X)$ is trivial, if n is even, and it is isomorphic to $[\mathbb{Z}/2\mathbb{Z}]^4$, if n is odd.

The group $\mathrm{Aut}_0(X)$ acts faithfully on the total cohomology ring $H^*(X, \mathbb{Z})$, when X is deformation equivalent to a generalized Kummer manifold, by a result of Oguiso [Og, Theorem 1.3]. An alternative proof of Theorem 1.1 for this deformation type follows from Oguiso's result via the argument used in the proof of Lemma 2.2 below.

The classification of irreducible holomorphic symplectic manifolds is an open problem. Two additional deformation types are known at present, one of six dimensional manifolds [O'G2] and one of ten dimensional manifolds [O'G1].

This work was motivated by the talk of Zhiyuan Li at the workshop "Hyper-Kähler Manifolds, Hodge Theory, and Chow Groups" at the Tsinghua Sanya International Mathematics Forum in December 2016. In his talk Li surveyed consequences of the existence of universal families over Teichmüller spaces to the study of cycles on moduli spaces of polarized irreducible holomorphic symplectic manifolds, generalizing previous work in the $K3$ surface case [BZMM].

2. Moduli spaces of marked pairs and triples

Fix a lattice Λ isometric to the Beauville-Bogomolov-Fujiki lattice of some irreducible holomorphic symplectic manifold. Set $\Lambda_{\mathbb{C}} := \Lambda \otimes_{\mathbb{Z}} \mathbb{C}$. Let Ω_{Λ} be the period domain of irreducible

holomorphic symplectic manifolds with Beauville-Bogomolov-Fujiki lattice Λ

$$\Omega_\Lambda := \{\ell \in \mathbb{P}(\Lambda_{\mathbb{C}}) \ : \ (\ell,\ell) = 0, \text{ and } (\ell,\bar{\ell}) > 0\}. \tag{2.1}$$

Fix a connected component $\mathfrak{M}^0_\Lambda$ of the moduli space of marked irreducible holomorphic symplectic manifolds. The *period map* $P : \mathfrak{M}^0_\Lambda \to \Omega_\Lambda$ sends the isomorphism class of a marked pair (X,η) to $\eta(H^{2,0}(X))$. Given a point $\ell \in \Omega_\Lambda$, denote by $\Lambda^{1,1}(\ell)$ the sublattice of Λ orthogonal to ℓ. The period map is a surjective local homeomorphism [Hu1]. P is generically injective and any two points in the same fiber of P are inseparable, by Verbitsky's Global Torelli Theorem [Ver, Hu2]. A point (X,η) in $\mathfrak{M}^0_\Lambda$ is a separated point, if and only if the Kähler cone of X is equal to its *positive cone*, where the latter is the connected component of $\{\alpha \in H^{1,1}(X,\mathbb{R}) \ : \ (\alpha,\alpha) > 0\}$ containing the Kähler cone [Ver, Hu2] (see also [Ma, Theorem 2.2 (4)]). Consequently, if $\Lambda^{1,1}(\ell)$ is trivial, or cyclic spanned by a class λ with $(\lambda,\lambda) \geq 0$, then the fiber $P^{-1}(\ell)$ consists of a single separated point of $\mathfrak{M}^0_\Lambda$ [Hu1, Corollaries 5.7 and 7.2] (see also [Ma, Theorem 2.2 (5)]).

Lemma 2.1. *Every local system over $\mathfrak{M}^0_\Lambda$ is trivial.*

Proof. The period domain Ω_Λ is simply-connected, by [Hu3, Cor. 3]. Hence, it suffices to prove that every local system $\mathcal{L}$ over $\mathfrak{M}^0_\Lambda$ is the pullback of a local system over Ω_Λ via the period map P. Choose a covering $\mathcal{U} := \{U_i \ : \ i \in I\}$ of $\mathfrak{M}^0_\Lambda$ by simply connected open subsets U_i, such that P restricts to each U_i as a homeomorphism. Set $V_i := P(U_i)$. We get the open covering $\mathcal{V} := \{V_i \ : \ i \in I\}$ of Ω_Λ, by the surjectivity of the period map. Given an n-tuple $\vec{i} := (i_1, i_2, \ldots, i_n) \in I^n$, denote by $J_{\vec{i}}$ the set of connected components of $U_{\vec{i}} := \cap_{k=1}^n U_{i_k}$. Set $V_{\vec{i}} := \cap_{k=1}^n V_{i_k}$. The period map restricts to an open embedding of $U_{\vec{i}}$ into $V_{\vec{i}}$. The complement of $P(U_{\vec{i}})$ in $V_{\vec{i}}$ is contained in the intersection of $V_{\vec{i}}$ with the countable union of closed complex analytic hyperplanes $\Omega_\Lambda \cap \lambda^\perp$, as λ varies in the set of primitive classes in Λ with $(\lambda,\lambda) < 0$, by the Global Torelli Theorem. Consequently, each connected component of $V_{\vec{i}}$ contains the image of a unique connected component of $U_{\vec{i}}$, by [Ver, Lemma 4.10]. We get a one-to-one correspondence between the set of connected components of $U_{\vec{i}}$ and $V_{\vec{i}}$.

The restriction $\mathcal{L}_{|_{U_i}}$ of $\mathcal{L}$ to U_i is a trivial local system, for all $i \in I$, as U_i is simply connected. Set $\Gamma(U_i,\mathcal{L}) := H^0(U_i, \mathcal{L}_{|_{U_i}})$. The evaluation homomorphism $ev^i_t : \Gamma(U_i,\mathcal{L}) \to \mathcal{L}_t$ is an isomorphism, for every fiber $\mathcal{L}_t$ of $\mathcal{L}$ over a point t of U_i. Given $\vec{i} \in I^2$ and $c \in J_{\vec{i}}$, denote by U_c the corresponding connected component of $U_{\vec{i}}$. Given points t_i and t'_i in U_i, parallel transport along any path from t_i to t'_i in U_i is given by $ev^i_{t'_i}(ev^i_{t_i})^{-1}$. The restriction homomorphism $\rho^{i_k}_c : \Gamma(U_{i_k},\mathcal{L}) \to \Gamma(U_c,\mathcal{L})$ is an isomorphism, for $k = 1,2$, and for every $c \in J_{\vec{i}}$. Hence, the gluing of $\mathcal{L}_{|_{U_{i_1}}}$ and $\mathcal{L}_{|_{U_{i_2}}}$ along U_c is determined by the isomorphism

$$f_{c,i_1,i_2} := (\rho^{i_1}_c)^{-1} \circ \rho^{i_2}_c : \Gamma(U_{i_2},\mathcal{L}) \to \Gamma(U_{i_1},\mathcal{L})$$

in the sense that the following diagram is commutative for a point t_c in U_c.



Given a point $t \in U_{i_1} \cap U_{i_2}$, set $g_{i_1,i_2}(t) := f_{c,i_1,i_2} : \Gamma(U_{i_2}, \mathcal{L}) \to \Gamma(U_{i_1}, \mathcal{L})$, where U_c is the connected component containing t. The transformation g_{i_1,i_2} glues the trivializations $\mathcal{L}_{|_{U_{i_1}}} \cong \underline{\Gamma(U_{i_1}, \mathcal{L})}_{U_{i_1}}$ and $\mathcal{L}_{|_{U_{i_2}}} \cong \underline{\Gamma(U_{i_2}, \mathcal{L})}_{U_{i_2}}$ along U_{i_1,i_2}.

Let $P_i : U_i \to V_i$ be the restriction of P to U_i. Denote by $P_i^* : \Gamma(V_i, P_{i,*}\mathcal{L}_{|_{U_i}}) \to \Gamma(U_i, \mathcal{L}_{|_{U_i}})$ the natural isomorphism and let $P_{i,*}$ be its inverse. We get the isomorphisms

$$\bar{f}_{c,i_1,i_2} := P_{i_1,*} \circ f_{c,i_1,i_2} \circ P_{i_2}^* : \Gamma(V_{i_2}, P_{i_2,*}\mathcal{L}_{|_{U_{i_2}}}) \to \Gamma(V_{i_1}, P_{i_1,*}\mathcal{L}_{|_{U_{i_1}}}).$$

The latter determines a gluing of $P_{i_1,*}\mathcal{L}_{|_{U_{i_1}}}$ and $P_{i_2,*}\mathcal{L}_{|_{U_{i_2}}}$ along the connected component V_c of V_{i_1,i_2}, for every $c \in J_{\vec{i}}$, hence along all the connected components of V_{i_1,i_2}. Denote by $\bar{g}_{i_1,i_2}$ the gluing of the trivializations $P_{i_k,*}\mathcal{L}_{|_{U_{i_k}}} \cong \underline{\Gamma(V_{i_k}, P_{i_k,*}\mathcal{L}_{|_{U_{i_k}}})}_{V_{i_k}}$, $k = 1, 2$, along V_{i_1,i_2}. These gluing transformations satisfy the co-cycle condition $\bar{g}_{i,k} = \bar{g}_{i,j}\bar{g}_{j,k}$, since each $\bar{g}_{i,j}$ pulls back to $g_{i,j}$ and the $g_{i,j}$'s satisfy the co-cycle condition. Let $\overline{\mathcal{L}}$ be the local system over Ω_Λ determined by the covering $\mathcal{V}$ and the gluing transformations $\bar{g}_{i_1,i_2}$. Then the restriction of $P^*\overline{\mathcal{L}}$ to U_i is naturally identified with that of $\mathcal{L}$ and each gluing transformations $P^*\bar{g}_{i_1,i_2}$ restricts to g_{i_1,i_2}. Hence, $\mathcal{L}$ is isomorphic to $P^*\overline{\mathcal{L}}$. □

We keep the notation of the introduction. Fix a group G isomorphic to the group $\mathrm{Aut}_0(X)$ of some irreducible holomorphic symplectic manifold X. The moduli space $\mathfrak{M}_{\Lambda,G}$ is constructed by gluing all Kuranishi families $\pi : \mathcal{X} \to U$, each endowed with a choice of an isometric trivialization $\eta : R^2\pi_*\mathbb{Z} \to \underline{\Lambda}_U$ and a trivialization $\psi : \mathrm{Aut}_0(\pi) \to \underline{G}_U$. The construction is completely analogous to that of $\mathfrak{M}_\Lambda$ in [Hu2, Prop. 4.3]. The following is an immediate corollary of Lemma 2.1.

Lemma 2.2. *The forgetful morphism $\phi : \mathfrak{M}_{\Lambda,G} \to \mathfrak{M}_\Lambda$ restricts to each connected component $\mathfrak{M}^0_{\Lambda,G}$ of $\mathfrak{M}_{\Lambda,G}$ as an isomorphism onto the corresponding connected component $\mathfrak{M}^0_\Lambda$ of $\mathfrak{M}_\Lambda$. The set of connected components of $\mathfrak{M}_{\Lambda,G}$ over $\mathfrak{M}^0_\Lambda$ forms a torsor under the group of outer automorphisms of G.*

Proof. Let $Ad_g \in \mathrm{Aut}(G)$ be conjugation by $g \in G$. G acts on the set $\mathrm{Isom}(\mathrm{Aut}_0(X), G)$, of group isomorphisms from $\mathrm{Aut}_0(X)$ onto G, by the action $\psi \mapsto Ad_g \circ \psi$. Similarly, $f \in \mathrm{Aut}_0(X)$ acts on $\mathrm{Isom}(\mathrm{Aut}_0(X), G)$ via $\psi \mapsto \psi \circ Ad_{f^{-1}}$. The set of orbits for the two actions coincide and we get a natural identification $\mathrm{Isom}(\mathrm{Aut}_0(X), G)/G = \mathrm{Isom}(\mathrm{Aut}_0(X), G)/\mathrm{Aut}_0(X)$, so we denote both orbit spaces by $\mathrm{Out}(\mathrm{Aut}_0(X), G)$. Over $\mathfrak{M}_\Lambda$ we have the local system $\mathcal{O}ut_G$, whose fiber over the point corresponding to the isomorphism class (X, η) is $\mathrm{Out}(\mathrm{Aut}_0(X), G)$. If f is an automorphism of a marked pair (X, η), then f belongs to $\mathrm{Aut}_0(X)$. Hence, the local system $\mathcal{O}ut_G$ is well defined.

The moduli space $\mathfrak{M}_{\Lambda,G}$ is simply the total space of the local system $\mathcal{O}ut_G$. The local system $\mathcal{O}ut_G$ restricts to a trivial local system over each connected component $\mathfrak{M}^0_\Lambda$, by Lemma 2.1. Hence, connected components of $\mathfrak{M}_{\Lambda,G}$ are simply global sections of $\mathcal{O}ut_G$. □

Given an abelian group A and a topological space S, denote by $\check{H}^i(S, A)$ the i-th Čech cohomology of S with coefficients in A.

Lemma 2.3. *The pull back homomorphism $P^* : \check{H}^i(\Omega_\Lambda, A) \to \check{H}^i(\mathfrak{M}^0_\Lambda, A)$ is an isomorphism, for every abelian group A.*

Proof. We keep the notation of the proof of Lemma 2.1. The simply connected open sets U of $\mathfrak{M}^0_\Lambda$, such that P restricts to U as a homeomorphism, form a basis for the topology of $\mathfrak{M}^0_\Lambda$.

Given an open covering $\mathcal{U} = \{U_i\}_{i\in I}$, consisting of such open sets, we get an open covering $\mathcal{V} := \{V_i := P(U_i)\}_{i\in I}$ of Ω_Λ. The covering $\mathcal{U}$ is a refinement of the covering $\{P^{-1}(V_i)\}_{i\in I}$, so we get a pullback homomorphism

$$P^* : \check{H}^i(\mathcal{V}, A) \to \check{H}^i(\mathcal{U}, A).$$

Every open covering of Ω_Λ admits a refinement by an open covering $\mathcal{V}$ as above. Hence, it suffices to prove that the pullback homomorphism displayed above is an isomorphism, for such a covering $\mathcal{U}$ of $\mathfrak{M}^0_\Lambda$ and the induced covering $\mathcal{V}$ of Ω_Λ. The degree n group $\mathcal{C}^n(\mathcal{U}, A)$ in the Čech complex is $\oplus_{\vec{i}\in I^{n+1}} \oplus_{c\in J_{\vec{i}}} \Gamma(U_c, A)$ and $\Gamma(U_c, A) = A$. The natural bijection between the connected components of $U_{\vec{i}}$ and $V_{\vec{i}}$, observed in the proof of Lemma 2.1, implies that the pullback homomorphism P^* induces an isomorphism of Čech complexes. □

3. Gerbes

Definition 3.1. A *fibered category* (or a *presheaf of categories*) $\mathbb{F}$ over a topological space S consists of the following.

(1) A category $\mathbb{F}(U)$ for each open subset U of X.
(2) A functor $i^* : \mathbb{F}(U) \to \mathbb{F}(V)$, for each inclusion $i : V \to U$ of open sets.
(3) A natural isomorphism $\tau_{ij} : (ij)^* \to j^*i^*$ for each composition $W \xrightarrow{j} V \xrightarrow{i} U$ of inclusions.

The natural isomorphisms are assumed to satisfy a natural associativity property [Gi, Br, Def. 5.2.1]. See also [Moe, Def. 2.1].

Definition 3.2. A fibered category $\mathbb{F}$ over a topological space S is called a *prestack* if, for any pair of objects a, b of $\mathbb{F}(U)$, the presheaf $\mathcal{H}om(a, b)$ over U, associating to an inclusion $i : V \to U$ of an open subset V the set $\mathrm{Hom}_{\mathbb{F}(V)}(i^*a, i^*b)$, is a sheaf. A prestack $\mathbb{F}$ is called a *stack*, if it satisfies an additional descent condition for objects [Gi, Br] (see also [Moe, Def. 2.6]).

Definition 3.3. ([Gi], [Br, Def. 5.2.4], or [Moe, Def. 3.1]). A *gerbe* over a topological space S is a stack $\mathbb{G}$ satisfying the following properties.

(1) Each category $\mathbb{G}(U)$ is a groupoid.
(2) S admits a covering by open sets U, such that $\mathbb{G}(U)$ has an object.
(3) Given objects a, b of $\mathbb{G}(U)$, any point $x \in U$ has an open neighborhood $V \subset U$, such that $\mathrm{Hom}_{\mathbb{G}(V)}(i^*a, i^*b)$ is non-empty, where $i : V \to U$ is the inclusion.

Let $\mathbb{G}$ be a gerbe over a topological space S, such that for every open set U and every object a of $\mathbb{G}(U)$, the group $\mathrm{Aut}_{\mathbb{G}(U)}(a)$ is abelian. We define next a sheaf $\mathcal{Z}$ of abelian groups called the *band* of $\mathbb{G}$. For a more general definition, for arbitrary gerbes, see [Gi] or [Moe, Def. 3.2]. We can choose an open covering $\mathcal{U} := \{U_\alpha\}$ and an object a_α in $\mathbb{G}(U_\alpha)$, yielding a sheaf $\mathcal{A}ut(a_\alpha)$ over each U_α, by Axiom (2) of Definition 3.3. These sheaves admit a canonical gluing to a sheaf $\mathcal{Z}$ of abelian groups as follows. Choose an covering $\{U^\xi_{\alpha\beta}\}$ of each $U_{\alpha\beta}$, such that there exists an isomorphism $f^\xi_{\alpha\beta}$ from the restriction $(a_\beta)_{|_{U^\xi_{\alpha\beta}}}$ of a_β to $U^\xi_{\alpha\beta}$ to the restriction $(a_\alpha)_{|_{U^\xi_{\alpha\beta}}}$ of a_α. Such a covering exists, by Axioms (1) and (3) of Definition 3.3. Conjugation by $f^\xi_{\alpha\beta}$ induces a sheaf isomorphism

$$\lambda^\xi_{\alpha\beta} : \mathcal{A}ut\left((a_\beta)_{|_{U^\xi_{\alpha\beta}}}\right) \to \mathcal{A}ut\left((a_\alpha)_{|_{U^\xi_{\alpha\beta}}}\right). \tag{3.1}$$

Now $\lambda^{\xi}_{\alpha\beta}$ is independent of the choice of $f^{\xi}_{\alpha\beta}$, since any other choice is of the form $hf^{\xi}_{\alpha\beta}$, for an element h of the abelian group $\mathrm{Aut}_{\mathbb{G}(U^{\xi}_{\alpha\beta})}\left((a_\alpha)_{|_{U^{\xi}_{\alpha\beta}}}\right)$. In particular, $\lambda^{\xi}_{\alpha\beta}$ and $\lambda^{\tilde{\xi}}_{\alpha\beta}$ agree over overlaps $U^{\xi}_{\alpha\beta} \cap U^{\tilde{\xi}}_{\alpha\beta}$ and define an isomorphism

$$\lambda_{\alpha\beta} : \mathcal{A}ut\left((a_\beta)_{|_{U_{\alpha\beta}}}\right) \to \mathcal{A}ut\left((a_\alpha)_{|_{U_{\alpha\beta}}}\right).$$

The isomorphisms $\lambda_{\alpha\beta}$ satisfy the co-cycle condition $\lambda_{\alpha\beta}\lambda_{\beta\gamma}\lambda_{\gamma\alpha} = 1$ in $\mathrm{Aut}\left(\mathcal{A}ut\left((a_\alpha)_{|_{U_{\alpha\beta\gamma}}}\right)\right)$, as the left hand side is an inner automorphism of an abelian group. Hence, the sheaves $\mathcal{A}ut(a_\alpha)$ glue to a sheaf $\mathcal{Z}$ of abelian groups, called the *band* of the gerbe $\mathbb{G}$. The isomorphism class of $\mathcal{Z}$ is independent of the choice of the covering $\{U_\alpha\}$ and of the objects $\{a_\alpha\}$.

Fix a lattice Λ and a finite group G. Let B be an analytic space and $\kappa : B \to \mathfrak{M}_{\Lambda,G}$ a morphism. Consider the fibered category $\kappa^{-1}\mathbb{G}$ over B, which associates to an open set $U \subset B$ the following category $\kappa^{-1}\mathbb{G}(U)$. Objects of $\kappa^{-1}\mathbb{G}(U)$ are triples (π, η, ψ) consisting of a family $\pi : \mathcal{X} \to U$ of irreducible holomorphic symplectic manifolds, an isometric trivialization $\eta : R^2\pi_*\mathbb{Z} \to \underline{\Lambda}_U$, and a trivialization $\psi : \mathcal{A}ut_0(\pi) \to \underline{G}_U$, such that for each point $u \in U$ the triple (X_u, η_u, ψ_u) represents the isomorphism class corresponding to the point $\kappa(u)$ of $\mathfrak{M}_{\Lambda,G}$. A morphism in $\mathrm{Hom}_{\kappa^{-1}\mathbb{G}(U)}((\pi,\eta,\psi),(\tilde{\pi},\tilde{\eta},\tilde{\psi}))$ is an isomorphism $f : \mathcal{X} \to \tilde{\mathcal{X}}$, satisfying $\pi = \tilde{\pi}f$, such that the induced isomorphism of local systems $f^* : R^2\tilde{\pi}_*\mathbb{Z} \to R^2\pi_*\mathbb{Z}$ satisfies $\tilde{\eta} = \eta f^*$ and such that $\tilde{\psi} = \psi Ad_f$, where $Ad_f : \mathcal{A}ut_0(\tilde{\pi}) \to \mathcal{A}ut_0(\pi)$ is the isomorphism of sheaves of groups induced by conjugation by f.

Lemma 3.4. *The fibered category $\kappa^{-1}\mathbb{G}$ is a gerbe over B with band $\underline{Z}_B$, where Z is the center of G.*

Proof. It suffices to prove that the fibered category $\mathbb{G}$ over $\mathfrak{M}_{\Lambda,G}$, associated to the identity morphism, is a gerbe with band $\underline{Z}$, as the more general $\kappa^{-1}\mathbb{G}$ described above is simply the inverse image of $\mathbb{G}$ via κ and is thus a gerbe with band $\underline{Z}_B$, by [Br, Prop. 5.2.6]. Property (1) of Definition 3.3 holds, by definition of morphisms in $\mathbb{G}$. The vector space $H^0(X, TX)$ vanishes for every irreducible holomorphic symplectic manifold X, and the dimension of $H^1(X, TX)$ is the Hodge number $h^{1,1}(X)$. Hence, the versal Kuranishi family of X is universal [BHPV, Ch. I, Theorems (10.3) and (10.5)]. Every point $t := (X_0, \eta_0)$ of $\mathfrak{M}_\Lambda$ admits a simply connected open neighborhood U, over which we have a universal family of deformations of X_0, as $\mathfrak{M}_\Lambda$ is constructed by gluing such families [Hu2, Prop. 4.3]. The same holds for $\mathfrak{M}_{\Lambda,G}$, by Lemma 2.2. Let b be a point of $\mathfrak{M}_{\Lambda,G}$ and let (X_b, η_b, ψ_b) represent the isomorphism class b. Choose a simply connected open neighborhood $U \subset \mathfrak{M}_{\Lambda,G}$ of b, over which we have a universal family $\pi : \mathcal{X} \to U$ of deformations of X_b. The local systems $R^2\pi_*\mathbb{Z}$ and $\mathcal{A}ut_0(\pi)$ are trivial, since U is simply connected, and so η_b and ψ_b extend to an isometric trivialization $\eta : R^2\pi_*\mathbb{Z} \to \underline{\Lambda}_U$ and a trivialization $\psi : \mathcal{A}ut_0(\pi) \to \underline{G}_U$. We get the object (π, η, ψ) in $\mathbb{G}(U)$. Property (2) of Definition 3.3 follows. Property (3) of Definition 3.3 holds, since the universal family is universal for each of its fibers (see [BHPV, Ch. I, Theorems (10.3) and (10.6) and the Remark following (10.6)]).

Given an open subset U of $\mathfrak{M}_{\Lambda,G}$ and an object $a := (\pi, \eta, \psi)$ in $\mathbb{G}(U)$, the sheaf $\mathcal{A}ut_{\mathbb{G}(U)}(a)$ is isomorphic to the center of $\mathcal{A}ut_0(\pi)$ and ψ restricts to an isomorphism from the center of $\mathcal{A}ut_0(\pi)$ onto $\underline{Z}_U$. The isomorphisms induced by the ψ's are compatible with the gluing transformations $\lambda^{\xi}_{\alpha\beta}$ in (3.1), by definition of morphisms in $\mathbb{G}$. Hence, the band of $\mathbb{G}$ is isomorphic to $\underline{Z}$. □

4. Vanishing of the class of a gerbe

Fix a lattice Λ isometric to the Beauville-Bogomolov-Fujiki lattice of some irreducible holomorphic symplectic manifold. Let W be a three dimensional positive definite subspace of $\Lambda_{\mathbb{R}}$. Let $Q_W \subset \mathbb{P}(W_{\mathbb{C}})$ be the conic of isotropic lines in $W_{\mathbb{C}} := W \otimes_{\mathbb{R}} \mathbb{C}$. Denote by $\iota : Q_W \to \Omega_\Lambda$ the inclusion into the period domain (2.1).

Lemma 4.1. *The inclusion ι is a homotopy equivalence.*

Proof. Let $Gr_+(3, \Lambda_{\mathbb{R}})$ be the Grassmannian of positive definite three dimensional subspaces of $\Lambda_{\mathbb{R}}$. The identity component $SO_+(\Lambda_{\mathbb{R}})$ of the special orthogonal group acts transitively on $Gr_+(3, \Lambda_{\mathbb{R}})$. The stabilizer of W in $SO_+(\Lambda_{\mathbb{R}})$ is $SO(W) \times SO(W^\perp)$ realizing $Gr_+(3, \Lambda_{\mathbb{R}})$ as the quotient $SO_+(\Lambda_{\mathbb{R}})/[SO(W) \times SO(W^\perp)]$. Now $SO(W) \times SO(W^\perp)$ is a maximal compact subgroup of $SO_+(\Lambda_{\mathbb{R}})$ and so the latter is topologically the product of the former and a Euclidean space, by Cartan's Theorem [Mos, Theorem 2]. Hence, $Gr_+(3, \Lambda_{\mathbb{R}})$ is contractible.

A point of the period domain Ω_Λ corresponds to an isotropic line $\ell \subset \Lambda_{\mathbb{C}}$, such that $[\ell + \bar{\ell}] \cap \Lambda_{\mathbb{R}}$ is a positive definite two-dimensional subspace V of $\Lambda_{\mathbb{R}}$. The natural identification of the two dimensional real vector spaces ℓ and V endows the latter with an orientation. This construction identifies the period domain Ω_Λ with the Grassmannian of oriented positive definite two-dimensional subspaces of $\Lambda_{\mathbb{R}}$ [Hu3]. Let $I \subset \Omega_\Lambda \times Gr_+(3, \Lambda_{\mathbb{R}})$ be the incidence correspondence. The fiber of the projection $q : I \to Gr_+(3, \Lambda_{\mathbb{R}})$ over W is Q_W. The inclusion $\tilde{\iota} : Q_W \to I$ of the fiber is a homotopy equivalence, since q is a fibration over a contractible base. The fiber of the projection $p : I \to \Omega_\Lambda$ over an oriented two dimensional subspace $V \subset \Lambda_{\mathbb{R}}$ is the projectivization of the positive cone in the subspace $V^\perp$ of signature $(1, b_2 - 3)$, i.e., a hyperbolic space. Hence, p is a homotopy equivalence, being a fibration with contractible fibers. We conclude that $\iota = p \circ \tilde{\iota}$ is the composition of two homotopy equivalences and so is such as well. □

Let X be an irreducible holomorphic symplectic manifold and ω a Kähler class on X. Set $V := [H^{2,0}(X) \oplus H^{0,2}(X)] \cap H^2(X, \mathbb{R})$ and set $W := V + \mathbb{R}\omega$. Then W is a positive definite three dimensional subspace of $H^2(X, \mathbb{R})$. There exists a unique Ricci flat hermetian metric g on X, whose imaginary part is a Kähler form representing the class ω, by Yau's proof of the Calabi conjecture [Be1]. Furthermore, there exists two additional complex structures I and J, covariantly constant with respect to the Riemannian metric which is the real part of the hermetian metric g, such that $IJ = K$. The identity, I, J, and K span a subalgebra of endomorphisms of the real tangent bundle, which is isomorphic to the algebra $\mathbb{H}$ of quaternions. The two-sphere

$$Tw_\omega := \{aI + bJ + cK \ : \ a^2 + b^2 + c^2 = 1\},$$

of purely imaginary unit quaternions, consists of integrable complex structures. The Riemannian metric and each of these complex structures $I_t \in Tw_\omega$ determine a Kähler form ω_t on the manifold $\overline{X}$ underlying X, hence a Hodge structure. Denote by X_t the complex manifold $(\overline{X}, I_t)$. The map $\kappa : Tw_\omega \to \mathbb{P}(H^2(\overline{X}, \mathbb{C}))$, sending I_t to $H^{2,0}(X_t)$, is a diffeomorphic embedding whose image is Q_W [Hu1, 1.17]. Endow Tw_ω with the complex structure of Q_W. We get a complex structure on the smooth manifold $\mathcal{X} := \overline{X} \times Tw_\omega$, such that the projection

$$\pi : \mathcal{X} \to Tw_\omega \tag{4.1}$$

is holomorphic, and the fiber of π over $I_t \in Tw_\omega$ is isomorphic to $(\overline{X}, I_t)$ [HKLR]. The above family is known as the *twistor family* associated to the Kähler form ω.

Choose a marking $\eta_0 : H^2(X,\mathbb{Z}) \to \Lambda$. It extends uniquely to an isometric trivialization $\eta : R^2\pi_*\mathbb{Z} \to \underline{\Lambda}_{Tw_\omega}$, since Tw_ω is simply connected. We get an embedding

$$\kappa_{\pi,\eta} : Tw_\omega \to \mathfrak{M}^0_\Lambda$$

into the connected component of $\mathfrak{M}_\Lambda$ containing (X,η_0).

Lemma 4.2. *The pullback homomorphism $\kappa^*_{\pi,\eta} : \check{H}^*(\mathfrak{M}^0_\Lambda, Z) \to \check{H}^*(Tw_\omega, Z)$ is an isomorphism, for every abelian group Z.*

Proof. The composition $P \circ \kappa_{\pi,\eta} : Tw_\omega \to \Omega_\Lambda$ with the period map P is the embedding of the base Tw_ω of the twistor family as the conic $Q_{\eta(W)}$ of isotropic lines in the complexification of the positive three dimensional subspace $\eta(W)$ of $\Lambda_\mathbb{R}$. Set $\iota := P \circ \kappa_{\pi,\eta}$. The pullback $\iota^* : \check{H}^*(\Omega_\Lambda, Z) \to \check{H}^2(Tw_\omega, Z)$ is an isomorphism, by Lemma 4.1, and the pullback $P^* : \check{H}^*(\mathfrak{M}^0_\Lambda, Z) \to \check{H}^*(\Omega_\Lambda, Z)$ is an isomorphism, by Lemma 2.3. Hence, $\kappa^*_{\pi,\eta} = \iota^* \circ (P^*)^{-1}$ is an isomorphism as well. □

Proof of Theorem 1.1. Choose some twistor family (4.1). The local system $\mathcal{A}ut_0(\pi)$ over Tw_ω is trivial, as the latter is simply connected. Choose a trivialization $\psi : \mathcal{A}ut_0(\pi) \to \underline{G}_{Tw_\omega}$, let $\kappa_{\pi,\eta,\psi} : Tw_\omega \to \mathfrak{M}_{\Lambda,G}$ be the classifying morphism, and let $\mathfrak{M}^0_{\Lambda,G}$ be the connected component containing its image. Let Z be the center of G. Then $\kappa^*_{\pi,\eta,\psi} : \check{H}^*(\mathfrak{M}^0_{\Lambda,G}, Z) \to \check{H}^*(Tw_\omega, Z)$ is an isomorphism, by Lemmas 4.2 and 2.2.

Let $\mathbb{G}$ be the gerbe over $\mathfrak{M}^0_{\Lambda,G}$ associated to the identity morphism from $\mathfrak{M}^0_{\Lambda,G}$ to itself as in Lemma 3.4. Let $\mathbb{F}$ be the gerbe over Tw_ω associated to the morphism $\kappa_{\pi,\eta,\psi}$ as in Lemma 3.4. Equivalence classes of gerbes over a space S with band a sheaf $\mathcal{Z}$ of abelian groups are in bijection with cohomology classes in $\check{H}^2(S,\mathcal{Z})$ [Gi, Br, Theorem 5.2.8]. The equivalence class $[\mathbb{F}]$ of $\mathbb{F}$ in $\check{H}^2(Tw_\omega, Z)$ vanishes, since we have the object (π,η,π) in $\mathbb{F}(Tw_\omega)$. The morphism $\kappa_{\pi,\eta,\psi}$ pulls back the class $[\mathbb{G}]$ of $\mathbb{G}$ in $\check{H}^2(\mathfrak{M}^0_{\Lambda,G}, Z)$ to $[\mathbb{F}]$. Hence, $\kappa^*_{\pi,\eta,\psi}([\mathbb{G}])$ vanishes. We conclude that the class $[\mathbb{G}]$ vanishes, since $\kappa^*_{\pi,\eta,\psi}$ is an isomorphism. Consequently, $\mathbb{G}(\mathfrak{M}^0_{\Lambda,G})$ has an object.

The set of isomorphism classes of objects of $\mathbb{G}(\mathfrak{M}^0_{\Lambda,G})$ is a torsor for the group $\check{H}^1(\mathfrak{M}^0_{\Lambda,G}, Z)$, by [Br, Prop. 5.2.5]. The latter is the trivial group, by Lemmas 4.2 and 2.2. Hence, $\mathbb{G}(\mathfrak{M}^0_{\Lambda,G})$ has a unique object, up to isomorphism. The forgetful morphism ϕ restricts to an isomorphism $\phi : \mathfrak{M}^0_{\Lambda,G} \to \mathfrak{M}^0_\Lambda$ (Lemma 2.2), transferring the object of $\mathbb{G}(\mathfrak{M}^0_{\Lambda,G})$ to a Λ-marked family $(\pi : \mathcal{X} \to \mathfrak{M}^0_\Lambda, \eta)$ over $\mathfrak{M}^0_\Lambda$.

It remains to prove the universal property of (π,η). Let $(\tilde{\pi} : \tilde{\mathcal{X}} \to B, \tilde{\eta})$ be a Λ-marked family over a connected analytic space B, let $\kappa : B \to \mathfrak{M}_\Lambda$ be the classifying morphism, and let $\mathfrak{M}^0_\Lambda$ be the connected component containing its image. The family $\pi : \mathcal{X} \to \mathfrak{M}^0_\Lambda$ is locally universal, as it restricts to a universal Kuranishi family over an open neighborhood of each point of $\mathfrak{M}^0_\Lambda$. Hence, the Λ-marked families $(\tilde{\pi},\tilde{\eta})$ and $(\kappa^*(\pi),\kappa^*(\eta))$ are locally isomorphic. Let $\mathcal{P} := \mathcal{I}som((\tilde{\pi},\tilde{\eta}),(\kappa^*(\pi),\kappa^*(\eta)))$ be the local system of isomorphisms of the two families, which are compatible with the markings. Then $\mathcal{P}$ is an $\mathcal{A}ut_0(\tilde{\pi})$ torsor and $(\kappa^*(\pi),\kappa^*(\eta))$ is isomorphic to $(\tilde{\pi},\tilde{\eta}) \times \mathcal{P}$. □

Remark 4.3. Let X be an irreducible holomorphic symplectic manifold, $\mathrm{Diff}(X)$ its diffeomorphism group, and $\mathrm{Diff}_0(X)$ the subgroup of elements isotopic to the identity. An easy sufficient criterion for the existence of a universal family over the connected component of Teichmüller space is the triviality of the intersection $\mathrm{Aut}(X) \cap \mathrm{Diff}_0(X)$ for every X in this connected component [C, Sec. 1.4]. Such X is called *rigidified* [C, Def. 12].

Remark 4.4. Let $(\tilde{\pi} : \tilde{\mathcal{X}} \to B, \tilde{\eta})$ be a Λ-marked family over a connected analytic space B, let $\kappa : B \to \mathfrak{M}^0_\Lambda$ be the classifying morphism, and let $(\kappa^*(\pi) : \kappa^*\mathcal{X} \to B, \kappa^*(\eta))$ be the pullback of the universal family. We can choose a trivialization $\psi : \mathcal{A}ut_0(\kappa^*(\pi)) \to \underline{G}_B$, by Lemma 2.2. Thus, the orbit of the isomorphism class of the principal G-bundle $\mathcal{P}_{(\tilde{\pi},\tilde{\eta})} := \mathcal{I}som((\kappa^*(\pi), \kappa^*(\eta)), (\tilde{\pi}, \tilde{\eta}))$, under the group $\mathrm{Out}(G)$ of outer automorphisms, is an invariant of the Λ-marked family $(\tilde{\pi}, \tilde{\eta})$. Let us reconstruct this invariant more directly under the assumption that the fibers of $\tilde{\pi}$ are rigidified (Remark 4.3). Fix a fiber X of $\tilde{\pi}$. Let $\mathrm{Diff}'(X)$ be the subgroup of $\mathrm{Diff}(X)$ fixing the connected component of Teichmüller space containing X and acting trivially on $H^2(X, \mathbb{Z})$. $\Gamma := \mathrm{Diff}'(X)/\mathrm{Diff}_0(X)$ is a subgroup of the mapping class group $\mathrm{Diff}(X)/\mathrm{Diff}_0(X)$. The natural homomorphism $h : \mathrm{Aut}_0(X) \to \Gamma$ is an isomorphism. Indeed, h is injective, as X is assumed to be rigidified, and surjective by [Ver, Theorem 4.26(iii) and Cor. 4.31] (the groups Γ and $\mathrm{Aut}_0(X)$ are denoted by G_I and K_I in [Ver]). The Λ-marked family $(\tilde{\pi}, \tilde{\eta})$ is differentiably locally trivial. Hence, there exists an open covering $\{U_i\}$ of B and trivializations $\tilde{X}_{|_{U_i}} \cong X \times U_i$, such that the gluing transformations are given by continuous maps from $U_i \cap U_j$ to $\mathrm{Diff}'(X)$. These gluing transformations yield a principal $\mathrm{Diff}'(X)$-bundle, hence a principal Γ-bundle, hence a principal $\mathrm{Aut}_0(X)$-bundle.

Acknowledgements: This work was partially supported by a grant from the Simons Foundation (#427110). I thank Zhiyuan Li for his interesting talk, and for asking whether universal families exist over Teichmüller spaces of holomorphic symplectic manifolds, at the workshop "Hyper-Kähler Manifolds, Hodge Theory, and Chow Groups" at the Tsinghua Sanya International Mathematics Forum in December 2016. I thank the organizers, Radu Laza, Kieran O'Grady, and Claire Voisin for the invitation to this interesting and instructive workshop. I thank Daniel Huybrechts and Sukhendu Mehrotra for helpful communications and for their comments on an earlier draft of this note.

References

[Be1] Beauville, A.: *Varietes Kähleriennes dont la premiere classe de Chern est nulle.* J. Diff. Geom. 18, p. 755–782 (1983).

[Be2] Beauville, A.: *Some remarks on Kähler manifolds with $c_1 = 0$.* Classification of algebraic and analytic manifolds (Katata, 1982), 1–26, Progr. Math., Birkhäuser Boston, Boston MA 1983.

[BHPV] Barth, W., Hulek, K., Peters, C., Van de Ven, A.: *Compact complex surfaces.* Second edition. Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics, 4. Springer-Verlag, Berlin, 2004.

[Br] Brylinski, J.: *Loop spaces, characteristic classes and geometric quantization.* Reprint of the 1993 edition. Modern Birkhäuser Classics. Birkhäuser Boston, Inc., Boston, MA, 2008.

[BSN] Boissière, S., Nieper-Wisskirchen, M., Sarti, A.: *Higher dimensional Enriques varieties and automiorphisms of generalized Kummer varieties.* J. Math. Pures Appl. 95 (2011) 553–563.

[BZMM] Bergeron, N., Li, Z., Millson, J., and Moeglin, C.: *The Noether-Lefschetz conjecture and generalizations.* Electronic preprint arXiv:1412.3774.

[C] Catanese, F.: *A superficial working guide to deformations and moduli. Handbook of moduli.* Vol. I, 161–215, Adv. Lect. Math. (ALM), 24, Int. Press, Somerville, MA, 2013.

[Gi] Giraud, J.: *Cohomologie non abélienne.* Die Grundlehren der mathematischen Wissenschaften, Band 179. Springer-Verlag, Berlin-New York, 1971.

[HKLR] Hitchin, N., Karlhede, A., Lindström, U., Roček, M.: *Hyper-Kähler metrics and supersymmetry.* Comm. Math. Phys. 108 (1987), no. 4, 535–589.

[HT] Hassett, B., Tschinkel, Y.: *Hodge theory and Lagrangian planes in generalized Kummer fourfolds.* Mosc. Math. J. 13 (2013), no. 1, 33–56.

[Hu1] Huybrechts, D.: *Compact Hyperkähler Manifolds: Basic results.* Invent. Math. 135 (1999), no. 1, 63-113 and Erratum: Invent. Math. 152 (2003), no. 1, 209–212.

[Hu2] Huybrechts, D.: *A global Torelli theorem for hyperkähler manifolds [after M. Verbitsky].* Séminaire Bourbaki: Vol. 2010/2011. Exposés 1027?1042. Astérisque No. 348 (2012), Exp. No. 1040, 375–403.

[Hu3] Huybrechts, D.: *Moduli spaces of hyperkähler manifolds and mirror symmetry.* Intersection theory and moduli, 185–247, ICTP Lect. Notes, XIX, Abdus Salam Int. Cent. Theoret. Phys., Trieste, 2004.

[Ma] Markman, E.: *A survey of Torelli and monodromy results for hyperkahler manifolds.* In "Complex and Differential Geometry", W. Ebeling et. al. (eds.), Springer Proceedings in Math. 8, (2011), 257–323.

[Moe] Moerdijk, I.: *Introduction to the language of stacks and gerbes.* Electronic preprint arXiv:0212266v1.

[Mos] Mostow, G. D.: *A new proof of E. Cartan's theorem on the topology of semi-simple groups.* Bull. Amer. Math. Soc. 55 (1949), 969–980.

[O'G1] O'Grady, K.: *Desingularized moduli spaces of sheaves on a $K3$.* J. Reine Angew. Math. 512 (1999), 49–117.

[O'G2] O'Grady, K.: *A new six-dimensional irreducible symplectic variety.* J. Algebraic Geom. 12 (2003), no. 3, 435–505.

[Og] Oguiso, K.: *No cohomologically trivial non-trivial automorphisms of generalized Kummer manifolds.* Electronic preprint arXiv:1208.3750v3.

[Ver] Verbitsky, M.: *Mapping class group and a global Torelli theorem for hyperkähler manifolds.* Duke Math. J. 162 (2013), no. 15, 2929–2986.

Department of Mathematics and Statistics, University of Massachusetts, Amherst, MA 01003, USA

E-mail address: markman@math.umass.edu